

File SM

National Aeronautics and Space Administration
Goddard Space Flight Center
Contract No. NAS-5-9299

ST - AI - 10508

RATES OF BASIC ION-MOLECULAR PROCESSES
IN THE IONOSPHERE

by

A. D. Danilov

(USSR)

GPO PRICE \$ _____

CFSTI PRICE(S) \$ _____

Hard copy (HC) 1.00

Microfiche (MF) 50

ff 653 July 65

N 67 12916

FACILITY FORM 802

(ACCESSION NUMBER)	(THRU)
<u>7</u>	<u>1</u>
(PAGES)	(CODE)
<u>CP-80151</u>	<u>13</u>
(NASA CR OR TMX OR AD NUMBER)	(CATEGORY)

17 AUGUST 1966

RATES OF BASIC ION-MOLECULAR PROCESSES
IN THE IONOSPHERE

Doklady A. N. SSSR, Geofizika,
Tom 169, No. 2, 332-4,
Izdatel'stvo "NAUKA," 1966

by A. D. Danilov


S U M M A R Y

The rate of basic ion-molecular processes are discussed on the basis of the review of twenty-nine works by various authors and the comparison of laboratory experiments with theoretical considerations.

*
* *

The determination of constants of the rate of ionization recombination processes in the upper atmosphere is beset with well known difficulties. As follows from the review paper [1], the span of laboratory investigations of the basic reaction constants is very broad and it does not allow to pinpoint reliable values for the investigation of ion-molecular reactions in the ionosphere and in the laboratory, which require discussion in order to select the most reliable values of constants.

We compiled in Table 1 the results of laboratory investigations of two basic ionospheric ion-molecular reactions



As may be seen from Table 1, even if we reject the high value of $2 \cdot 10^{-10} \text{ cm}^3 \text{ sec}^{-1}$, brought out by Paulson (inasmuch as it is in contradiction with the estimate [9], $\gamma_1 \leq 5 \cdot 10^{-11} \text{ cm}^3 \text{ sec}^{-1}$ brought out in the same author's review paper), and also the high value $\gamma_1 \approx 10^{-8} \text{ cm}^3 \cdot \text{sec}^{-1}$, which,

TABLE 1

No.	A U T H O R S	Constants of reactions (1) and (2) in $\text{cm}^3 \text{ sec}^{-1}$	
		γ_1	γ_2
1	R. F. Potter [2]	10^{-8}	-
2	R. H. D. Dickinson [3]	-	$2.5 \cdot 10^{-11}$
3	V. L. Talrose et al [4]	$\leq 6.7 \cdot 10^{-11}$	-
4	[5] G. F. O. Langstroth, J. B. Hast.	$4.7 \cdot 10^{-12}$	$1.8 \cdot 10^{-12}$
5	Fite, et al. [6]	-	$(1-10) \cdot 10^{-11}$
6	Volpi, et al. [7]	$2.2 \cdot 10^{-11}$	-
7	Sayers, Smith [8]	$2.7 \cdot 10^{-11}$	$1.6 \cdot 10^{-11}$
8	Paulson, Mosher	$\leq 5 \cdot 10^{-11}$	-
9	Paulson, et al. [10]	$2 \cdot 10^{-10}$	-
10	Fehsenfeld, et al. [11] [12]	$3 \cdot 10^{-12}$	$4 \cdot 10^{-11}$

as already noted more than once, is probably erroneous, there is observed a scattering of the most reliable among the obtained values of the constant of the reaction's (1) rate by one order of magnitude; at the same time, both the upper and the lower values are corroborated by two experiments. A somewhat better position is observed for the constant of reaction (2); however, here too it is necessary to deny oneself the result of the experiment [5], which as will be seen further, will give a value γ_1 more acceptable from the viewpoint of ionospheric investigations so that the remaining values γ_2 may be matched with a precision to the factor 2.

The estimates of the effectiveness of the same processes (1) and (2), obtained by various authors on the basis of analysis of ionospheric data, are compiled in Table 2. Here it should be noted that different initial data were utilized by the various authors, as for example: the altitude variation of the ion composition in the region 100–200 km [16], the altitude and daily variations of ion concentrations [17], the measurements of the linear recombination coefficient in the F₂-region [19] and so forth. In order to obtain the estimates brought out in the eighth line of Table 2, the relations between ion-molecular reaction and dissociative recombination constants [24–26]

$$\gamma_1 / \alpha^* \text{NO}^+ = (5-10) \cdot 10^{-5}, \quad \gamma_2 / \alpha^* \text{O}_2^+ = (1-2) \cdot 10^{-4}$$

were utilized. The quantities $\alpha^* \text{NO}^+$ and $\alpha^* \text{O}_2^+$, which were more reliably investigated in the laboratory than the constants γ_1 and γ_2 , were borrowed from the review paper [2] for a temperature of the order of 1000° K. Therefore, the estimates of γ_1 and γ_2 brought up above, also refer to the indicated temperature.

It may be seen from the data brought out in Table 2 that the ionospheric

TABLE 2

No.	A U T H O R S	Constants of Reaction (1) and (2) $\text{cm}^3 \text{sec}^{-1}$	
		γ_1	γ_2
1.	Bates, Nicolet [14]	$\gamma_1 + 0.16$	$2 = 1.3 \cdot 10^{-13}$
2.	Khartek, Rive [15]	$\leq 10^{-13}$	$\leq 10^{-12}$
3.	Norton, et al [16]	$1 \cdot 10^{-12}$	$5 \cdot 10^{-11}$
4.	Danilov [17]	$(0.5-1) \cdot 10^{-12}$	$(0.5-1) \cdot 10^{-11}$
5.	Whitten, Poppoff [18]	$2 \cdot 10^{-12}$	$2 \cdot 10^{-11}$
6.	Nisbet, Quinn [19]	$\leq 1.3 \cdot 10^{-12}$	$\leq 10^{-11}$
7.	Sagalyn, Smiddy [20]	$(1.6-3.2) \cdot 10^{-12}$	—
8.	Danilov (γ_1/α^* & α^*) [21]	$(1.5-3) \cdot 10^{-12}$	$(0.6-1.1) \cdot 10^{-11}$
9.	Hall, et al [22]	$\gamma_1 + 0.12$	$\gamma_2 \approx 3 \cdot 10^{-12}$
10.	Danilov, Yatsenko [23]	$\gamma_1/\gamma_2 \approx 0.1$	

estimates by the various authors are not generally divergent among themselves, leading to respective mean values of constants of processes (1) and (2)

$$\gamma_1 = 2 \cdot 10^{-12} \text{ cm}^3 \cdot \text{sec}^{-1}$$

$$\gamma_2 = 2 \cdot 10^{-11} \text{ cm}^3 \cdot \text{sec}^{-1}$$

these mean values of γ not differing from the most reliable estimates brought out in the lines 3 to 9 of Table 2 by more than 2—3 times. The values of γ , obtained in [14, 15] should be recognized as underrated. They agree with neither the most reliable ionospheric estimates, nor the laboratory data. As to the latter, the quantities γ_1 and γ_2 obtained from the ionospheric data allow to assert that the quantities γ_1 arrived at in the works [5, 11], are correct whereas those obtained in the experiments [7, 8] are overrated.

It should be analogously admitted that the value

$$\gamma_2 = 1.8 \cdot 10^{-12} \text{ cm}^3 \cdot \text{sec}^{-1}$$

is strongly underrated, while that obtained in recent experiments

$$\gamma_2 = 4 \cdot 10^{-11} \text{ cm}^3 \cdot \text{sec}^{-1}$$

is somewhat overrated.

The obtained estimates of constants of basic ion-molecular reactions in the ionosphere appear to be sufficiently reliable (in any case, within the limits of a factor of 2). Some uncertainty is introduced only by the temperature dependence of constants γ_1 and γ_2 . According to the experiments of the work [8], the reaction's (2) constant depends on the temperature in the form T^{-1} in the region $210^\circ \leq T \leq 452^\circ \text{ K}$. From theoretical considerations, it appears that generally the ion-molecular reaction constants' dependence on temperature must not be observed [9, 27, 28]. According to [13] a direct dependence between γ_1 and T was obtained in the experiments at energies of several eV. The question of the constants' temperature dependence thus requires further experimental research.

TABLE 3

Reactions	Fite [6] et al	Volpi [7] et al	Paulson & al [10]	Ferguson [11-13] & al	Harteck [29]
$N^+ + O_2 \rightarrow NO^+ + O$ (3)	$5 \cdot 10^{-10}$	$1 \cdot 10^{-10}$	$3 \cdot 10^{-10}$	$5 \cdot 10^{-10}$	10^{-10}
$O_2^+ + N_2 \rightarrow NO^+ + NO$ (4)	—	$\leq 2 \cdot 10^{-10}$	$4 \cdot 10^{-11}$	$< 10^{-10}$	—
$N_2^+ + O_2 \rightarrow NO^+ + NO$ (5)	$2,1 \cdot 10^{-13}$	$\leq 2 \cdot 10^{-10}$	$4 \cdot 10^{-11}$	—	—
$N_2^+ + O_2 \rightarrow O_2^+ + N_2$ (6)	$2 \cdot 10^{-10}$	—	—	$1,0 \cdot 10^{-10}$	—
$N_2^+ + O \rightarrow NO^+ + N$ (7)	—	—	—	$2,5 \cdot 10^{-10}$	—
$N_2^+ + O \rightarrow N_2 + O^+$ (8)	—	—	—	$< 10^{-11}$	—
$N^+ + O_2 \rightarrow O_2^+ + N$ (9)	—	—	—	$(0,5-1) \cdot 10^{-9}$	—

Compiled in Table are the results of laboratory measurements of constants of ion-molecular reactions in which the main components of the atmosphere O , O_2 and N_2 take part. The bulk of these reactions was investigated in the works [11-13]. As in Table 1, attention is drawn here by the values obtained by Paulson, sharply diverging from the data by other authors, which compels us to assume that the Paulson experiment [10] is erroneous. The ionospheric estimates of these reactions' constants are more complex than those of the basic reactions (1) and (2). Particular attention should be given the values

$$\gamma_7 \approx 1,0 \cdot 10^{-11} \text{ cm}^3 \text{ sec}^{-1}$$

obtained in [16]

$$\gamma_7 \approx 2 \cdot 10^{-11} \text{ cm}^3 \text{ sec}^{-1}$$

obtained in [18], and also

$$\gamma_4 = (3-10) \cdot 10^{-14} \text{ cm}^3 \text{ sec}^{-1}$$

made in the work [26] on the basis of analysis of data on the ion composition of the region 100-150 km.

The results of laboratory experiments on the constants of ion-molecular processes with the participation of small atmospheric admixtures, such as the neutral molecules NO and nitrogen atoms, are compiled in Table 4

TABLE 4

Reactions	Value of γ according to [13]
$O^+ + NO \rightarrow NO^+ + O$ (10)	$(2,4 \pm 1) \cdot 10^{-11}$
$N_2^+ + NO \rightarrow NO^+ + N_2$ (11)	$5 \cdot 10^{-10}$
$O_2^+ + NO \rightarrow NO^+ + O_2$ (12)	$8 \cdot 10^{-10}$
$O_2^+ + N \rightarrow NO^+ + O$ (13)	$(1,8 \pm 0,5) \cdot 10^{-10}$
$N^+ + NO \rightarrow NO^+ + N$ (14)	$(8 \pm 1) \cdot 10^{-10}$

As may be seen, the values of constants obtained in the laboratory are very high. Therefore, the opinion having prevailed to-date that the indicated reactions may be disregarded on account of low concentrations of NO and N must be revised. The exact values of the indicated concentrations are so far unknown; however, the high values of constants require detailed analysis

of these processes when considering the equilibrium concentrations of ions, particularly in the region 100–130 km.

*** THE END ***

Institute of Applied Physics,
Received on 20 October 1965
Contract No. NAS-5-9299
Consultants & Designers, Inc.,
Arlington, Va.

Translated by
ANDRE L. BRICHANT
on
18 August 1966

REFERENCES

1. A. D. DANILOV, G. S. IVANOV-KHOLODNYI. UFN, 85 (2), 259, 1960.
2. R. F. POTTER. J. Chem. Phys., 23, 2462, 1955.
3. P. H. D. DICKINSON, J. SAYERS. Proc. Phys. Soc., 76, 137, 1960.
4. V. L. TALROSE, et al. Disk. Farad. Soc., 33, 257, 1962.
5. G. F. O. LANGSTROTH, J. B. HASTED. Disk. Farad. Soc., 33, 298, 1962.
6. W. L. FITE, et al. Disk. Farad. Soc., 33, 246, 1962.
7. G. G. VOLPI, et al. J. Chem. Phys., 39, 518, 1963.
8. J. SAYERS, D. SMITH. Disk. Farad. Soc., 37, 112, 1964.
9. J. F. PAULSON. Symposium D'Aeronomie, p. 18, 1964.
10. J. F. PAULSON, et al. Proc. Nat. Acad. Sci., India, A, 33 (4), 522, 1963.
11. F. C. FEHSENFELD, et al. Planet. Sp. Sci., 13, 219, 1965.
12. F. C. FEHSENFELD, et al. Planet Sp. Sci., 13, 579, 1965.
13. E. E. FERGUSON. J. Geophys. Res., 70, No. 17, 4323, 1965.
14. D. R. BATES, M. NICOLET. J. Atmos. and Terr. Phys., 18, 65, 1960.
15. P. KHARTEK, R. RIVS. Sborn. Elementarnyye protsessy v verkhney atmosfere, M., str. 7, 1965.
16. R. B. NORTON, et al. Proc. Intern. Conf. Ionosphere, London, 1963.
17. A. D. DANILOV. Kosmicheskiye issledovaniya, 2 (6), 865, 1964.
18. R. C. WHITTEN, I. G. POPPOFF. J. Atmos. Sci., 21, 117, 1964.
19. J. S. NISBET, T. P. QUINN. J. Geophys. Res., 70 (1), 113, 1965.
20. R. C. SAGALYN, M. SMIDDY. J. Geophys. Res., 69, 1809, 1964.
21. G. S. IVANOV-KHOLODNYI. Geomagnetizm i aeronomiya, 6, No. 4, 1966.
22. L. A. HALL, et al. Space Research, 5, 1965.
23. A. D. DANILOV, S. P. YATSENKO. Kosmicheskiye issledovaniya, 2 (2), 276, 1964.
24. A. D. DANILOV. Iskusstvennyye sputniki Zemli, v. 5, 1960; v. 7, 1961.
25. G. S. IVANOV-KHOLODNYI. DAN, 137, No. 2, 327, 1961.
26. G. S. IVANOV-KHOLODNYI, A. D. DANILOV. Kosmicheskiye issledovaniya, 4, No. 3, 1966.
27. F. H. FIELD, et al. J. Am. Chem. Soc., 79, 2419, 1957.
28. H. EYRING, et al. J. Chem. Phys., 4, 479, 1936.
29. P. HARTECK. Disk. Farad. Soc., No. 37, 224, 1964.

D I S T R I B U T I O NGODDARD S. F. C.NASA HQS.OTHER CENTERS

100	CLARK, TOWNSEND	SS	NEWELL, NAUGLE	<u>AMES</u>
110	STROUD	SG	MITCHELL, SMITH	SONETT
400	BOURDEAU		SCHARDT	LIBRARY
601	FAVA		SCHMERLING	
610	MEREDITH		DUBIN	<u>LANGLEY</u>
	SEDDON	SC	LIDDEL	160 ADAMSON
611	MC DONALD	SL	FELLOWS	185 WEATHERWAX
	ABRAHAM		HIPSHER	
	BOLDT, VKB		HOROWITZ	<u>J. P. L.</u>
	SERLEMITSOS	SM	FOSTER, ALLENBY	32 MEGHREBLIAN
	TEEGARDEN		GILL	186-133
612	HEPPNER	RR	KURZWEG	412 WYCKOFF Vis-Lab
	NESS		RTR NEILL	752 LIBRARY
613	KUPPERIAN		USS-T	
	ALEXANDER			<u>U. of IOWA</u>
	REED			VAN ALLEN
	HALLAM			
614	WHITE			<u>U. of C. at BERKELEY</u>
	FROST			WILCOX
	KASTNER			
615	BAUER			<u>S. W. C. A. S.</u>
	AIKIN			JOHNSON
	GOLDBERG, HERMAN			
	MAIER, STONE			
	KANE			
	CHANDRA			
	DONLEY			
640	HESS (3)			
641	MEAD, MAEDA			
	HARRIS			
	HERRING			
620	SPENCER			
621	NEWTON			
	TAYLOR			
630	GI for SS (3)			
252	LIBRARY			
256	FREAS			
